UNIVERSITY OF HAWAII LIBRARY

Journal of MOLECULAR BIOLOGY

JUL 23 1985

EDITOR-IN-CHIEF
J. C. KENDREW

ASSISTANT EDITOR
S. BRENNER

EDITORIAL BOARD
C. R. Cantor
P. Chambon
W. W. Franke
M. Gellert
G. A. Gilbert
M. Gottesman
I. Herskowitz
R. Huber
H. E. Huxley
A. Klug
V. Luzzati
J. H. Miller
M. F. Moody
K. Simons

VOLUME 180

25 November to 25 December 1984

ICAL
H
01
73
BRARY

ACADEMIC PRESS
London New York San Francisco

Copyright © 1984, by Academic Press Inc. (London) Ltd.

ALL RIGHTS RESERVED

No part of this volume may be reproduced in any form, by photostat, microfilm, or any other means, without written permission from the publishers.

Printed in Great Britain

CONTENTS OF VOLUME 180

Number 1, 25 November 1984

Campbell, M. E. M., Palfreyman, J. W. and Preston, C. M., Identification of Herpes Simplex Virus DNA Sequences Which Encode a *Trans*-acting Polypeptide Responsible for Stimulation of Immediate Early Transcription 1

Shepherd, B. M. and Finnegan, D. J., Structure of Circular Copies of the *412* Transposable Element Present in *Drosophila melanogaster* Tissue Culture Cells, and Isolation of a Free *412* Long Terminal Repeat . . 21

Wemmer, D. E., Chou, S.-H. and Reid, B. R., Sequence-specific Recognition of DNA. Nuclear Magnetic Resonance Assignments and Structural Comparison of Wild-type and Mutant λ OR3 Operator DNA 41

Hahn, S., Dunn, T. and Schleif, R., Upstream Repression and CRP Stimulation of the *Escherichia coli* L-Arabinose Operon 61

Edwards, C. A. and Firtel, R. A., Site-specific Phasing in the Chromatin of the rDNA in *Dictyostelium discoideum* 73

Gwynne, D. I., Miller, B. L., Miller, K. Y. and Timberlake, W. E., Structure and Regulated Expression of the SpoCl Gene Cluster from *Aspergillus nidulans* 91

Graessmann, M., Guhl, E., Bumke-Vogt, C. and Graessmann, A., The Second Large Simian Virus 40 T-antigen Exon Contains the Information for Maximal Cell Transformation 111

Hannon, R., Bateman, E., Allan, J., Harborne, N. and Gould, H., Control of RNA Polymerase Binding to Chromatin by Variations in Linker Histone Composition 131

Van Stolk, B. J. and Noller, H. F., Chemical Probing of Conformation in Large RNA Molecules. Analysis of 16 S Ribosomal RNA using Diethylpyrocarbonate 151

Lilley, D. M. J. and Hallam, L. R., Thermodynamics of the ColEl Cruciform. Comparisons Between Probing and Topological Experiments Using Single Topoisomers 179

Letters to the Editor

Dunn, T. M. and Schleif, R., Deletion Analysis of the *Escherichia coli ara* P_C and P_{BAD} Promoters 201

Miller, J. H., Genetic Studies of the *lac* Repressor. XII. Amino Acid Replacements in the DNA Binding Domain of the *Escherichia coli lac* Repressor 205

Lebioda, L. and Brewer, J. M., Crystallization and Preliminary Crystallographic Data for a Tetragonal Form of Yeast Enolase . 213

Number 2, 5 December 1984

LeClerc, J. E., Istock, N. L., Saran, B. R. and Allen, R., Jr, Sequence Analysis of Ultraviolet-induced Mutations in M13*lacZ* Hybrid Phage DNA 217

Kammerer, B., Guyonvarch, A. and Hubert, J. C., Yeast Regulatory Gene *PPR1*. I. Nucleotide Sequence, Restriction Map and Codon Usage 239

Liljelund, P., Losson, R., Kammerer, B. and Lacroute, F., Yeast Regulatory Gene *PPR1*. II. Chromosomal Localization, Meiotic Map, Suppressibility, Dominance/Recessivity and Dosage Effect . . . 251

Lane, D., Hill, D., Caughey, P and Gunn, P., The mini-F Primary Origin. Sequence Analysis and Multiple Activities 267

Frackman, S., Siegele, D. A. and Feiss, M., A Functional Domain of Bacteriophage λ Terminase for Prohead Binding 283

Wlodawer, A., Walter, J., Huber, R. and Sjölin, L., Structure of Bovine Pancreatic Trypsin Inhibitor. Results of Joint Neutron and X-ray Refinement of Crystal Form II 301

Trinick, J., Knight, P. and Whiting, A., Purification and Properties of Native Titin 331

Lecomte, J., La Mar, G. N., Winterhalter, K. H. and Smit, J. D. G., Proton Nuclear Magnetic Resonance Investigation of the Heme Cavity Structure of Liver Fluke (*Dicrocoelium dendriticum*) Methemoglobin . 357

Letters to the Editor

Cheng, K. C. and Smith, G. R., Recombinational Hotspot Activity of Chi-like Sequences 371

Brandt, P. W., Diamond, M. S. and Schachat, F. H., The Thin Filament of Vertebrate Skeletal Muscle Co-operatively Activates as a Unit 379

Deisenhofer, J., Epp, O., Miki, K., Huber, R. and Michel, H., X-ray Structure Analysis of a Membrane Protein Complex. Electron Density Map at 3 Å Resolution and a Model of the Chromophores of the Photosynthetic Reaction Center from *Rhodopseudomonas viridis* 385

Number 3, 15 December 1985

Parker, M. L., Christensen, A. C., Boosman, A., Stockard, J., Young, E. T. and Doermann, A. H., Nucleotide Sequence of Bacteriophage T4 Gene *23* and the Amino Acid Sequence of its Product 399

KHANDEKAR, P., SAIDAPET, C., KRAUSKOPF, M., ZARRAGA, A.-M., LIN, W.-L., MENDOLA, C. and SIDDIQUI, M. A. Q., Co-ordinate Control of Gene Expression. Muscle-specific 7 S RNA Contains Sequences Homologous to 3′-Untranslated Regions of Myosin Genes and Repetitive DNA 417

O'HARE, K., MURPHY, C., LEVIS, R. and RUBIN, G. M., DNA Sequence of the *white* Locus of *Drosophila melanogaster* 437

QUAX-JEUKEN, Y., JANSSEN, C., QUAX, W., VAN DEN HEUVEL, R. and BLOEMENDAL, H., Bovine β-Crystallin Complementary DNA Clones. Alternating Proline/Alanine Sequence of βB1 Subunit Originates from a Repetitive DNA Sequence 457

WATERSTON, R. H., HIRSH, D. and LANE, T. R., Dominant Mutations Affecting Muscle Structure in *Caenorhabditis elegans* that Map near the Actin Gene Cluster 473

LANDEL, C. P., KRAUSE, M., WATERSTON, R. H. and HIRSH, D., DNA Rearrangements of the Actin Gene Cluster in *Caenorhabditis elegans* Accompany Reversion of Three Muscle Mutants 497

BEEN, M. D. and CHAMPOUX, J. J., Breakage of Single-stranded DNA by Eukaryotic Type 1 Topoisomerase Occurs Only at Regions with the Potential for Base-pairing 515

COSTANZO, M., BRZUSTOWICZ, L., HANNETT, N. and PERO, J., Bacteriophage SPO1 Genes *33* and *34*. Location and Primary Structure of Genes Encoding Regulatory Subunits of *Bacillus subtilis* RNA Polymerase . 533

VARENNE, S., BUC, J., LLOUBES, R. and LAZDUNSKI, C., Translation is a Non-uniform Process. Effect of tRNA Availability on the Rate of Elongation of Nascent Polypeptide Chains 549

SERGEANT, A., BOHMANN, D., ZENTGRAF, H., WEIHER, H. and KELLER, W., A Transcription Enhancer Acts *in Vitro* over Distances of Hundreds of Base-pairs on both Circular and Linear Templates but not on Chromatin-reconstituted DNA 577

KIROV, N., DJONDJUROV, L. and TSANEV, R., Nuclear Matrix and Transcriptional Activity of the Mouse α-Globin Gene 601

KREIG, P., STRACHAN, R., WALLIS, E., TABE, L. and COLMAN, A., Efficient Expression of Cloned Complementary DNAs for Secretory Proteins after Injection into *Xenopus* Oocytes 615

TABE, L., KRIEG, P., STRACHAN, R., JACKSON, D., WALLIS, E. and COLMAN, A., Segregation of Mutant Ovalbumins and Ovalbumin–Globin Fusion Proteins in *Xenopus* Oocytes 645

UENO, H. and HARRINGTON, W. F., An Enzyme–Probe Study of Motile Domains in the Subfragment-2 Region of Myosin 667

Luo, M., Arnold, E., Erickson, J. W., Rossmann, M. G., Boege, U. and Scraba, D. G., Picornaviruses of Two Different Genera have Similar Structures 703

Wüthrich, K., Billeter, M. and Braun, W., Polypeptide Secondary Structure Determination by Nuclear Magnetic Resonance Observation of Short Proton–Proton Distances 715

Appendix: Short Intramolecular ^1H–^1H Distances in Polypeptide Chains Containing Glycine Proline or *cis* Peptide Bonds. . . 734

Pardi, A., Billeter, M. and Wüthrich, K., Calibration of the Angular Dependence of the Amide Proton–C^α Proton Coupling Constants, $^3J_{HN\alpha}$, in a Globular Protein. Use of $^3J_{HN\alpha}$ for Identification of Helical Secondary Structure 741

Letters to the Editor

Lustig, A. J. and Petes, T. D., Long Poly(A) Tracts in the Human Genome are Associated with the *Alu* Family of Repeated Elements 753

Silverton, E. W., Padlan, E. A., Davies, D. R., Smith-Gill, S. and Potter, M., Crystalline Monoclonal Antibody Fabs Complexed to Hen Egg White Lysozyme 761

Number 4, 25 December 1985

Chang, L.-Y. E. and Slightom, J. L., Isolation and Nucleotide Sequence Analysis of the β-Type Globin Pseudogene from Human, Gorilla and Chimpanzee 767

Harris, S., Barrie, P. A., Weiss, M. L. and Jeffreys, A. J., The Primate $\psi\beta 1$ Gene. An Ancient β-Globin Pseudogene. 785

Goodman, M., Koop, B. F., Czelusniak, J., Weiss, M. L. and Slightom, J. L., The η-Globin Gene. Its Long Evolutionary History in the β-Globin Gene Family of Mammals 803

Alonso, A., Jorcano, J. L., Beck, E., Hovemann, B. and Schmidt, T., *Drosophila melanogaster* U1 snRNA Genes 825

Bryant, J. L. Jr and King, J., DNA Injection Proteins are Targets of Acridine-sensitized Photoinactivation of Bacteriophage P22 . . 837

Place, N., Fien, K., Mahoney, M. E., Wulff, D. L., Ho, Y.-S., Debouck, C., Rosenberg, M., Shih, M.-C. and Gussin, G. N., Mutations that Alter the DNA Binding Site for the Bacteriophage Lambda *c*II Protein and Affect the Translation Efficiency of the *c*II Gene 865

Malan, T. P., Kolb, A., Buc, H. and McClure, W. R., Mechanism of CRP-cAMP Activation of *lac* Operon Transcription Initiation Activation of the P1 Promoter 881

Appendix 900

Sass, H. and Pederson, T., Transcription-dependent Localization of U1 and U2 Small Nuclear Ribonucleoproteins at Major Sites of Gene Activity in Polytene Chromosomes 911

Myslinski, E., Branlant, C., Wieben, E. D. and Pederson, T., The Small Nuclear RNAs of *Drosophila* 927

Lin, W.-L. and Pederson, T., Ribonucleoprotein Organization of Eukaryotic RNA. XXXI. Structure of the U1 Small Nuclear Ribonucleoprotein 947

Weaver, D. T. and DePamphilis, M. L., The Role of Palindromic and Non-palindromic Sequences in Arresting DNA Synthesis *in Vitro* and *in Vivo* 961

Bozzoni, I., Fragapane, P., Annesi, F., Pierandrei-Amaldi, P., Amaldi, F. and Beccari, E., Expression of Two *Xenopus laevis* Ribosomal Protein Genes in Injected Frog Oocytes. A Specific Splicing Block Interferes with the L1 RNA Maturation 987

Khalili, K. and Weinmann, R., Actin mRNAs in HeLa Cells. Stabilization after Adenovirus Infection 1007

Hudson, G. S. and Davidson, B. E., Nucleotide Sequence and Transcription of the Phenylalanine and Tyrosine Operons of *Escherichia coli* K12 1023

Olson, E. R., Tomich, C.-S. C. and Friedman, D. I., The *nusA* Recognition Site. Alteration in Its Sequence or Position Relative to Upstream Translation Interferes with the Action of the N Anti-termination Function of Phage Lambda 1053

Adoutte, A., de Loubresse, N. G. and Beisson, J., Proteolytic Cleavage and Maturation of the Crystalline Secretion Products of *Paramecium* . 1065

Goodenough, U. and Heuser, J., Structural Comparison of Purified Dynein Proteins with *In Situ* Dynein Arms 1083

Moo-Penn, W. F., Jue, D. L., Johnson, M. H., McDonald, M. J., Turci, S. M., Shih, T.-B., Jones, R. T., Therrell, B. L. Jr and Arnone, A., Structural and Functional Studies of Hemoglobin Wayne: An Elongated α-Chain Variant 1119

Brems, D. N. and Baldwin, R. L., Amide Proton Exchange Used to Monitor the Formation of a Stable α-Helix by Residues 3 to 13 During Folding of Ribonuclease S 1141

Vitrano, E., Cupane, A. and Cordone, L., Oxygen Affinity of Bovine Haemoglobin. Relevance of Electrostatic and Hydrophobic Interactions 1157

Appendix 1167

LETTERS TO THE EDITOR

BOZZONI, I., ANNESI, F., BECCARI, E., FRAGAPANE, P., PIERANDREI-AMALDI, P. and AMALDI, F., Splicing of *Xenopus laevis* Ribosomal Protein RNAs is inhibited *in Vivo* by Antisera to Ribonucleoproteins Containing U1 Small Nuclear RNA 1173

TAKAGI, T., IIDA, S., MATSUOKA, A. and SHIKAMA, K., *Aplysia* Myoglobins with an Unusual Amino Acid Sequence 1179

ZANETTE, D., MONACO, H. L., ZANOTTI, G. and SPADON, P., Crystallization of Hen Eggwhite Riboflavin-binding Protein . . 1185

PARGE, H. E., SCHNEIDER, M., HAHN, U., SAENGER, W., ALTSCHMIED, L. and HILLEN, W., Crystallization of and Preliminary X-ray Diffraction Data for TET-repressor and the TET-repressor–Tetracycline Complex 1189

ERRATUM 1193

Index to Volume 180

Acridine-sensitized photoinactivation of bacteriophage P22; DNA injection proteins are targets, 837
Actin gene cluster, and nearby dominant mutations affecting muscle structure in *Caenorhabditis elegans*, 473
Actin gene cluster in *Caenorbaditis elegans*, DNA rearrangements accompany reversion of three muscle mutants, 497
Actin mRNAs in HeLa cells. Stabilization after adenovirus infection, 1007
Adenovirus-infected HeLa cells, actin mRNAs, 1007
Adoutte, A., De Loubresse, N. G. & Beisson, J., Proteolytic cleavage and maturation of the crystalline secretion products of *Paramecium*, 1065
Allan, J. (see also Hannon, R.), 131
Allen, R., Jr (see also Leclerc, J. E.), 217
Alonso, A., Jorcano, J. L., Beck, E., Hovemann, B. & Schmidt, T., *Drosophila melanogaster* U1 snRNA genes, 825
Altschmied, L. (see also Parge, H. E.), 1189
Alu family of repeated elements in associated with long poly(A) tracts in the human genome, 753
Amaldi, F. (see also Bozzoni, I.), 987, 1173
Amide proton exchange used to monitor the formation of a stable α-helix by residues 3 to 13 during folding of ribonuclease S, 1141
Amide proton–C^{α} proton coupling constants, $^{3}J_{HN\alpha}$, in a globular protein, calibration of the angular dependence; use for identification of helical secondary structure, 741
Amino acid replacements in the DNA binding domain of the *Escherichia coli lac* repressor, 205
Amino acid sequence of T4 gene *23* product, 399
Annesi, F. (see also Bozzoni, I.), 987, 1173
Antibody Fabs, monoclonal, crystalline, complexed to hen egg white lysozyme, 761
Antisera to ribonucleoproteins containing U1 small nuclear RNA inhibit *in vivo* the splicing of *Xenopus laevis* ribosomal protein RNAs, 1173
Aplysia myoglobins with an unusual amino acid sequence, 1179
ara P_C and P_{BAD} promoters, *Escherichia coli*; deletion analysis, 201
L-Arabinose operon, *Escherichia coli*, upstream and CRP stimulation, 61
Arnold, E. (see also Luo, M.), 703
Arnone, A. (see also McDonald, M. J.), 1119
Aspergillus nidulans, structure and regulated expression of the SpoC1 gene cluster, 91

Bacillus subtilis RNA polymerase, location and primary structure of genes encoding regulatory subunits; bacteriophage SP01 genes *33* and *34*, 533
Bacteriophage lambda, alteration in the sequence or position of the *nusA* recognition site interferes with the action of the N antitermination function, 1053
Bacteriophage lambda cII protein, mutations that alter the DNA binding site, 865
Bacteriophage λ OR3 operator DNA, wild-type and mutant, n.m.r. assignments and structural comparison, 41
Bacteriophage λ terminase, a functional domain for prohead binding, 283
Bacteriophage P22, DNA injection proteins are targets of acridine-sensitized photoinactivation, 837
Bacteriophage SP01 genese *33* and *34*. Location and primary structure of genes encoding regulatory subunits of *Bacillus subtilis* RNA polymerase, 533
Bacteriophage T4 gene *23* nucleotide sequence and the amino acid sequence of its product, 399
Baldwin, R. L. (see also Brems, D. N.), 1141
Barrie, P. A. (see also Harris, S.), 785
Bateman, E. (see also Hannon, R.), 131
Beccari, E. (see also Bozzoni, I.), 987, 1173
Beck, E. (see also Alonso, A.), 825
Been, M. D. & Champoux, J. J., Breakage of single-stranded DNA by eukaryotic type 1 topoisomerase occurs only at regions with the potential for base-pairing, 515
Beisson, J. (see also Adoutte, A.), 1065
Billeter, M. (see also Pardi, A.), 741
Billeter, M. (see also Wüthrich, K.), 715

LETTERS TO THE EDITOR

BOZZONI, I., ANNESI, F., BECCARI, E., FRAGAPANE, P., PIERANDREI-AMALDI, P. and AMALDI, F., Splicing of *Xenopus laevis* Ribosomal Protein RNAs is inhibited *in Vivo* by Antisera to Ribonucleoproteins Containing U1 Small Nuclear RNA 1173

TAKAGI, T., IIDA, S., MATSUOKA, A. and SHIKAMA, K., *Aplysia* Myoglobins with an Unusual Amino Acid Sequence . . . 1179

ZANETTE, D., MONACO, H. L., ZANOTTI, G. and SPADON, P., Crystallization of Hen Eggwhite Riboflavin-binding Protein . . 1185

PARGE, H. E., SCHNEIDER, M., HAHN, U., SAENGER, W., ALTSCHMIED, L. and HILLEN, W., Crystallization of and Preliminary X-ray Diffraction Data for TET-repressor and the TET-repressor–Tetracycline Complex 1189

ERRATUM 1193

Index to Volume 180

Acridine-sensitized photoinactivation of bacteriophage P22; DNA injection proteins are targets, 837
Actin gene cluster, and nearby dominant mutations affecting muscle structure in *Caenorhabditis elegans*, 473
Actin gene cluster in *Caenorbaditis elegans*, DNA rearrangements accompany reversion of three muscle mutants, 497
Actin mRNAs in HeLa cells. Stabilization after adenovirus infection, 1007
Adenovirus-infected HeLa cells, actin mRNAs, 1007
Adoutte, A., De Loubresse, N. G. & Beisson, J., Proteolytic cleavage and maturation of the crystalline secretion products of *Paramecium*, 1065
Allan, J. (see also Hannon, R.), 131
Allen, R., Jr (see also Leclerc, J. E.), 217
Alonso, A., Jorcano, J. L., Beck, E., Hovemann, B. & Schmidt, T., *Drosophila melanogaster* U1 snRNA genes, 825
Altschmied, L. (see also Parge, H. E.), 1189
Alu family of repeated elements in associated with long poly(A) tracts in the human genome, 753
Amaldi, F. (see also Bozzoni, I.), 987, 1173
Amide proton exchange used to monitor the formation of a stable α-helix by residues 3 to 13 during folding of ribonuclease S, 1141
Amide proton–C^{α} proton coupling constants, $^{3}J_{HN\alpha}$, in a globular protein, calibration of the angular dependence; use for identification of helical secondary structure, 741
Amino acid replacements in the DNA binding domain of the *Escherichia coli lac* repressor, 205
Amino acid sequence of T4 gene *23* product, 399
Annesi, F. (see also Bozzoni, I.), 987, 1173
Antibody Fabs, monoclonal, crystalline, complexed to hen egg white lysozyme, 761
Antisera to ribonucleoproteins containing U1 small nuclear RNA inhibit *in vivo* the splicing of *Xenopus laevis* ribosomal protein RNAs, 1173
Aplysia myoglobins with an unusual amino acid sequence, 1179
ara P_C and P_{BAD} promoters, *Escherichia coli*; deletion analysis, 201
L-Arabinose operon, *Escherichia coli*, upstream and CRP stimulation, 61
Arnold, E. (see also Luo, M.), 703
Arnone, A. (see also McDonald, M. J.), 1119
Aspergillus nidulans, structure and regulated expression of the SpoC1 gene cluster, 91

Bacillus subtilis RNA polymerase, location and primary structure of genes encoding regulatory subunits; bacteriophage SP01 genes *33* and *34*, 533
Bacteriophage lambda, alteration in the sequence or position of the *nusA* recognition site interferes with the action of the N antitermination function, 1053
Bacteriophage lambda cII protein, mutations that alter the DNA binding site, 865
Bacteriophage λ OR3 operator DNA, wild-type and mutant, n.m.r. assignments and structural comparison, 41
Bacteriophage λ terminase, a functional domain for prohead binding, 283
Bacteriophage P22, DNA injection proteins are targets of acridine-sensitized photoinactivation, 837
Bacteriophage SP01 genese *33* and *34*. Location and primary structure of genes encoding regulatory subunits of *Bacillus subtilis* RNA polymerase, 533
Bacteriophage T4 gene *23* nucleotide sequence and the amino acid sequence of its product, 399
Baldwin, R. L. (see also Brems, D. N.), 1141
Barrie, P. A. (see also Harris, S.), 785
Bateman, E. (see also Hannon, R.), 131
Beccari, E. (see also Bozzoni, I.), 987, 1173
Beck, E. (see also Alonso, A.), 825
Been, M. D. & Champoux, J. J., Breakage of single-stranded DNA by eukaryotic type 1 topoisomerase occurs only at regions with the potential for base-pairing, 515
Beisson, J. (see also Adoutte, A.), 1065
Billeter, M. (see also Pardi, A.), 741
Billeter, M. (see also Wüthrich, K.), 715

Bloemendal, H. (see also Quax-Jeuken, Y.), 457
Boege, U. (see also Luo, M.), 703
Bohmann, D. (see also Sergeant, A.), 577
Boosman, A. (see also Parker, M. L.), 399
Bovine β-crystallin complementary DNA clones. Alternating proline/alanine sequence of βB1 subunit originates from a repetitive DNA sequence, 457
Bovine haemoglobin oxygen affinity; relevance of electrostatic and hydrophobic interactions, 1157
Bovine pancreatic trypsin inhibitor structure; results of joint neutron and X-ray refinement of crystal form II, 301
Bozzoni, I., Annesi, F., Beccari, E., Fragapane, P., Pierandrei-Amaldi, P. & Amaldi, F., Splicing of *Xenopus laevis* ribosomal protein RNAs is inhibited *in vivo* by antisera to ribonucleoproteins containing U1 small nuclear RNA, 1173
Bozzoni, I., Fragapane, P., Annesi, F., Pierandrei-Amaldi, P., Amaldi, F. & Beccari, E., Expression of two *Xenopus laevis* ribosomal protein genes in injected frog oocytes. A specific splicing block interferes with the L1 RNA maturation, 987
Brandt, P. W., Diamond, M. S. & Schachat, F. H., The thin filament of vertebrate skeletal muscle co-operatively activates as a unit, 379
Branlant, C. (see also Myslinski, E.), 927
Braun, W. (see also Wüthrich, K.), 715
Breakage of single-stranded DNA by eukaryotic type 1 topoisomerase occurs only at regions with the potential for base-pairing, 515
Brems, D. N. & Baldwin, R. L., Amide proton exchange used to monitor the formation of a stable α-helix by residues 3 to 13 during folding of ribonuclease S, 1141
Brewer, J. M. (see also Lebioda, L.), 213
Bryant, J. L. Jr & King, J., DNA injection proteins are targets of acridine-sensitized photoinactivation of bacteriophage P22, 837
Brzustowicz, L. (see also Costanzo, M.), 533
Buc, H. (see also Malan, T. P.), 881
Buc, J. (see also Varenne, S.), 549
Bumke-Vogt, C. (see also Graessman, M.), 111

cII gene translation efficiency is affected by mutations that alter the DNA binding site for the bacteriophage lambda cII protein, 865
Caenorhabditis elegans, DNA rearrangements of the actin gene cluster accompany reversion of three muscle mutants, 497
Caenorhabditis elegans, dominant mutations affecting muscle structure that map near the actin gene cluster, 473
Campbell, M. E. M., Palfreyman, J. W. & Preston, C. M., Identification of herpes simplex virus DNA sequences which encode a *trans*-acting polypeptide responsible for stimulation of immediate early transcription, 1
Caughey, P. (see also Lane, D.), 267
Cell transformation, maximal; the information is contained in the second large SV40 T-antigen exon, 111
Champoux, J. J. (see also Been, M. D.), 515
Chang, L.-Y. E. & Slightom, J. L., Isolation and nucleotide sequence analysis of the β-type globin pseudogene from human, gorilla and chimpanzee, 767
Chemical probing of conformation in large RNA molecules. Analysis of 16 S ribosomal RNA using diethylpyrocarbonate, 151
Cheng, K. C. & Smith, G. R., Recombinational hotspot activity of Chi-like sequences, 371
Chi-like sequences, recombinational hotspot activity, 371
Chimpanzee, gorilla and human, isolation and nucleotide sequence analysis of the β-type globin pseudogene, 767
Chou, S.-H. (see also Wemmer, D. E.), 41
Christensen, A. C. (see also Parker, M. L.), 399
Chromatin, control of RNA polymerase binding by variations in linker histone composition, 131
Chromatin of the rDNA in *Dictyostelium discoideum*, site-specific phasing, 73
Chromatin-reconstituted DNA is not acted on by a transcription enhancer *in vitro*, 577
Chromophores of the photosynthetic reaction centre from *Rhodopseudomonas viridis*, a model from X-ray structure analysis, 385
Chromosomal localization, yeast regulatory gene *PPR1*, 251

Chromosomes, polytene, transcription dependent localization of U1 and U2 small nuclear ribonucleoproteins at major sites of gene activity, 911
cis peptide bonds, and short proton–proton distances in polypeptide chains, 734
Cloned complementary DNAs for secretory proteins, efficient expression after injection into *Xenopus* oocytes, 615
Clones, bovine β-crystallin complementary DNA. Alternating proline/alanine sequence of βB1 subunit originates from a repetitive DNA sequence, 457
Codon usage, yeast regulatory gene *PPR1*, 239
ColE1 cruciform thermodynamics. Comparisons between probing and topological experiments using single topoisomers, 179
Colman, A. (see also Krieg, P.), 615
Colman, A. (see also Tabe, L.), 645
Conformation, chemical probing in large RNA molecules, Analysis of 16 S ribosomal RNA using diethylpyrocarbonate, 151
Co-operative activation as a unit of the thin filament of vertebrate skeletal muscle, 379
Co-ordinate control of gene expression. Muscle-specific 7 S RNA contains sequences homologous to 3′-untranslated regions of myosin genes and repetitive DNA, 417
Cordone, L. (see also Vitrano, E.), 1157
Costanzo, M., Brzustowicz, L., Hannett, N. & Pero, J., Bacteriophage SP01 genes *33* and *34*. Location and primary structure of genes encoding regulatory subunits of *Bacillus subtilis* RNA polymerase, 533
CRP-cAMP mechanism of *lac* operon transcription initiation activation of the P1 promoter, 881
CRP stimulation of the *Escherichia coli* L-arabinose operon, 61
Crystal form II structure of bovine pancreatic trypsin inhibitor, neutron and X-ray refinement, 301
β-Crystallin complementary DNA clones, bovine. Alternating proline/alanine sequence of βB1 subunit originates from a repetitive DNA sequence, 457
Crystalline monoclonal antibody Fabs complexed to hen egg white lysozyme, 761
— secretion products of *Paramecium*, proteolytic cleavage and maturation, 1065
Crystallization of hen eggwhite riboflavin-binding protein, 1185
— and preliminary crystallographic data for a tetragonal form of yeast enolase, 213
— of and preliminary X-ray diffraction data for TET-repressor and the TET-repressor–tetracycline complex, 1189
Cupane, A. (see also Vitrano, E.), 1157
Czelusniak, J. (see also Goodman, M.), 803

Davidson, B. E. (see also Hudon, G. S.), 1023
Davies, D. R. (see also Silverton, E. W.), 761
Debouck, C. (see also Place, N.), 865
Deisenhofer, J., Epp, O., Miki, K., Huber, R. & Michel, H., X-ray structure analysis of a membrane protein complex. Electron density map at 3 Å resolution and a model of the chromophores of the photosynthetic reaction center from *Rhodopseudomonas viridis*, 385
Deletion analysis of the *Escherichia coli ara* P_C and P_{BAD} promoters, 201
DNA, chromatin-reconstituted, is not acted on by a transcription enhancer *in vitro*, 577
—, complementary, bovine β-crystallin, clones. Alternating proline/alanine sequence of βB1 subunit originates from a repetitive DNA sequence, 457
— binding domain of the *Escherichia coli lac* repressor, amino acid replacements, 205
— — site for the bacteriophage lambda cII protein, altering mutations, 865
— injection proteins are targets of acridine-sensitized photoinactivation of bacteriophage P22, 837
—, M13*lacZ* hybrid phage, sequence analysis of ultraviolet-induced mutations, 217
— rearrangements of the actin gene cluster in *Caenorhabditis elegans* accompany reversion of three muscle mutants, 497
—, repetitive, muscle-specific 7 S RNA contains sequences homologous to 3′-untranslated regions; co-ordinate control of gene expression, 417
—, ribosomal, in *Dictyostelium discoideum*, site-specific phasing in the chromatin, 73
— sequence of the *white* locus of *Drosophila melanogaster*, 437
— sequences, herpes simplex virus, which encode a *trans*-acting polypeptide responsible for stimulation of immediate early transcription, 1
— sequence-specific recognition; n.m.r. assignments and structural comparison of wild-type and mutant λ OR3 operator DNA, 41
DNA, single-stranded, breakage by eukaryotic type 1 topoisomerase occurs only at regions with the potential for base-pairing, 515

DNA synthesis arrest, the role of palindromic and non-palindromic sequences, *in vitro* and *in vivo*, 961
DNAs, complementary, cloned, for secretory proteins, efficient expression after injection into *Xenopus* oocytes, 615
DePamphilis, M. L. (see also Weaver, D. T.), 961
Diamond, M. S. (see also Brandt, P. W.), 379
Dicrocoelium dendriticum methemoglobin, proton n.m.r. investigation of the heme cavity structure, 357
Dictyostelium discoideum, site-specific phasing in the chromatin of the rDNA, 73
Diethylpyrocarbonate probing of conformation of 16 S ribosomal RNA, 151
Djondjurov, L. (see also Kirov, N.), 601
Doermann, A. H. (see also Parker, M. L.), 399
Dominance/recessivity, yeast regulatory gene *PPR1*, 251
Dosage effect, yeast regulatory gene *PPR1*, 251
Drosophila melanogaster, DNA sequence of the *white* locus, 437
— —, tissue culture cells, structure of circular copies of the *412* transposable element; and isolation of a free *412* long terminal repeat, 21
Drosophila, the small nuclear RNAs, 927
— *melanogaster* U1 snRNA genes, 825
Dunn, T. (see also Hahn, S.), 61
Dunn, T. M. & Schleif, R., Deletion analysis of the *Escherichia coli ara* P_C and P_{BAD} promoters, 201
Dynein proteins, purified, structural comparison with *in situ* dynein arms, 1083

Edwards, C. A. & Firtel, R. A., Site-specific phasing in the chromatin of the rDNA in *Dictyostelium discoideum*, 73
Enolase, yeast, crystallization and preliminary crystallographic data for a tetragonal form, 213
Enzyme-probe study of motile domains in the subfragment-2 region of myosin, 667
Epp, O. (see also Deisenhofer, J.), 385
Erickson, J. W. (see also Luo, M.), 703
Escherichia coli ara P_C and P_{BAD} promoters, deletion analysis, 201
— — K12, nucleotide sequence and transcription of the phenylalanine and tyrosine operons, 1023
— — L-arabinose operon, upstream repression and CRP stimulation, 61
— — *lac* repressor, amino acid replacements in the DNA binding domain, 205
Eukaryotic RNA, structure of the U1 small nuclear ribonucleoprotein, 947
— type 1 topoisomerase breakage of single-stranded DNA occurs only at regions with the potential for base-pairing, 515
Evolutionary history of the η-globin gene in the β-globin gene family of mammals, 803
Expression of cloned complementary DNAs for secretory proteins after injection into *Xenopus* oocytes, 615
—, regulated, of the SpoC1 gene cluster from *Aspergillus nidulans*, 91
- of two *Xenopus laevis* ribosomal protein genes in injected frog oocytes. A specific splicing block interferes with the L1 RNA maturation, 987

Fabs, crystalline monoclonal antibody, complexed to hen egg white lysozyme, 761
Feiss, M. (see also Frackman, S.), 283
Fien, K. (see also Place, N.), 865
Filament, thin, of vertebrate skeletal muscle co-operatively activates as a unit, 379
Finnegan, D. J. (see also Shepherd, B. M.), 21
Firtel, R. A. (see also Edwards, C. A.), 73
Folding of ribonuclease S, and amide proton exchange used to monitor the formation of a stable α-helix by residues 3 to 13, 1141
412 transposable element in *Drosophila melanogaster* tissue culture cells, structure of circular copies; and isolation of a free *412* long terminal repeat, 21
Frackman, S., Siegele, D. A. & Feiss, M., A functional domain of bacteriophage λ terminase for prohead binding, 283
Fragapane, P. (see also Bozzoni, I.), 987, 1173
Friedman, D. I. (see also Olson, E. R.), 1053
Frog oocytes, injected, and the expression of two *Xenopus laevis* ribosomal protein genes; a specific block interferes with the L1 RNA maturation, 987
Functional domain of bacteriophage λ terminase for prohead binding, 283

Gene *c*II translation efficiency is affected by mutations that alter the DNA binding site for the bacteriophage lambda *c*II protein, 865
— cluster, actin, in *Caenorhabditis elegans*, DNA rearrangements accompany reversion of three muscle mutants, 497
— —, —, and nearby dominant mutations affecting muscle structure in *Caenorhabditis elegans*, 473
— — SpoC1 from *Aspergillus nidulans*, structure and regulated expression, 91
—, η-globin; its long evolutionary history in the β-globin gene family of mammals, 803
— expression co-ordinate control. Muscle-specific 7 S RNA contains sequences homologous to 3′-untranslated regions of myosin genes and repetitive DNA, 417
—, mouse α-globin; nuclear matrix and transcriptional activity, 601
— *PPR1*, yeast; chromosomal localization, meiotic map, suppressibility, dominance/recessivity and dosage effect, 251
— —, —; nucleotide sequence, restriction map and codon usage, 239
— $\psi\beta$1, primate; an ancient β-globin pseudogene, 785
— *23*, T4, nucleotide sequence, and the amino acid sequence of its product, 399
Genes, ribosomal protein, *Xenopus laevis*, two, expression in injected frog oocytes; a specific splicing block interferes with the L1 RNA maturation, 987
— snRNA, U1, *Drosophila melanogaster*, 825
— *33* and *34*, bacteriophage SP01; location and primary structure of genes encoding regulatory subunits of *Bacillus subtilis* RNA polymerase, 533
Genetic studies of the *lac* repressor. Amino acid replacements in the DNA binding domain of the *Escherichia coli lac* repressor, 205
Genome, human, long poly(A) tracts are associated with the *Alu* family of repeated elements, 753
α-Globin gene, mouse; nuclear matrix and transcriptional activity, 601
β-Globin gene family of mammals, and the long evolutionary history of the η-globin gene, 803
— pseudogene, ancient; the primate $\psi\beta$1 gene, 785
— — from human, gorilla and chimpanzee, isolation and nucleotide sequence analysis, 767
η-Globin gene; its long evolutionary history in the β-globin gene family of mammals, 803
Globular protein, calibration of the angular dependence of the amide proton–C^{α} coupling constants, $^{3}J_{HN\alpha}$; use for identification of helical secondary structure, 741
Glycine peptide bonds, and short proton-proton distances in polypeptide chains, 734
Goodenough, U. & Heuser, J., Structural comparison of purified dynein proteins with *in situ* dynein arms, 1083
Goodman, M., Koop, B. F., Czelusniak, J., Weiss, M. L. & Slightom, J. L., The η-globin gene. Its long evolutionary history in the β-globin gene family of mammals, 803
Gorilla, chimpanzee and human, isolation and nucleotide sequence analysis of the β-type globin pseudogene, 767
Gould, H. (see also Hannon, R.), 131
Graessmann, A. (see also Graessmann, M.), 111
—, M., Guhl, E., Bumke-Vogt, C. & Graessmann, A., The second large simian virus 40 T-antigen exon contains the information for maximal cell transformation, 111
Guhl, E. (see also Graessman, M.), 111
Gunn, P. (see also Lane, D.), 267
Gussin, G. N. (see also Place, N.), 865
Guyonvarch, A. (see also Kammerer, B.), 239
Gwynne, D. I., Miller, B. L., Miller, K. Y. & Timberlake, W. E., Structure and regulated expression of the SpoC1 gene cluster from *Aspergillus nidulans*, 91

Haemoglobin, bovine, oxygen affinity; relevance of electrostatic and hydrophobic interactions, 1157
Hahn, S., Dunn, T. & Schleif, R., Upstream repression and CRP stimulation of the *Escherichia coli* L-arabinose operon, 61
—, U. (see also Parge, H. E.), 1189
Hallam, L. R. (see also Lilley, D. M. J.), 179
Hannett, N. (see also Costanzo, M.), 533
Hannon, R., Bateman, E., Allan, J., Harborne. N. & Gould, H., Control of RNA polymerase binding to chromatin by variations in linker histone composition, 131
Harborne, N. (see also Hannon, R.), 131
Harrington, W. F. (see also Ueno, H.), 667
Harris, S., Barrier, P. A., Weiss, M. L. & Jeffreys, A. J., The primate $\psi\beta$1 gene. An ancient β-globin pseudogene, 785
HeLa cells, actin mRNAs; stabilization after adenovirus infection, 1007

α-Helix formation by residues 3 to 13 during folding of ribonuclease S, use of amide proton exchange for monitoring, 1141
Heme cavity structure of liver fluke methemoglobin, proton n.m.r. investigation, 357
Hemoglobin Wayne: an elongated α-chain variant; structural and functional studies, 1119
Hen egg white lysozyme complexes with crystalline monoclonal antibody Fabs, 761
— — — riboflavin-binding protein, crystallization, 1185
Herpes simplex virus DNA sequences which encode a *trans*-acting polypeptide responsible for stimulation of immediate early transcription, 1
Heuser, J. (see also Goodenough, U.), 1083
Hill, D. (see also Lane, D.), 267
Hillen, W. (see also Parge, H. E.), 1189
Hirsh, D. (see also Landel, C. P.), 497
Hirsh, D. (see also Waterson, R. H.), 473
Histone, linker, composition variations control RNA polymerase binding to chromatin, 131
Ho, Y.-S. (see also Place, N.), 865
Hotspot recombinational activity of Chi-like sequences, 371
Hovemann, B. (see also Alonso, A.), 825
Huber, R. (see also Deisenhofer, J.), 385
—, —. — — Wlodawer, A.), 301
Hubert, J. C. (see also Kammerer, B.), 239
Hudson, G. S. & Davidson, B. E., Nucleotide sequence and transcription of the phenylalanine and tyrosine operons of *Escherichia coli* K12, 1023
Human genome, long poly(A) tracts are associated with the *Alu* family of repeated elements, 753
—, gorilla and chimpanzee, isolation and nucleotide sequence analysis of the β-type globin pseudogene, 767
Hybrid phage M13*lacZ* DNA, sequence analysis of ultraviolet-induced mutations, 217
Hydrophobic and electrostatic interactions, relevance to oxygen affinity of bovine haemoglobin, 1157

Iida, S. (see also Takagi, T.), 1179
Istock, N. L. (see also Leclerc, J. E.), 217

$^3J_{HN\alpha}$, amide proton–C^α proton coupling constants, in a globular protein, calibration of the angular dependence; use for identification of helical secondary structure, 741
Jackson, D. (see also Tabe, L.), 645
Janssen, C. (see also Quax-Jeuken, Y.), 457
Jeffreys, A. J. (see also Harris, S.), 785
Johnson, M. H. (see also Moo-Penn, W. F.), 119
Jones, R. T. (see also Moo-Penn, W. F.), 1119
Jorcano, J. L. (see also Alonso, A.), 825
Jue, D. L. (see also Moo-Penn, W. F.), 1119

Kammerer, B., Guyonvarch, A. & Hubert, J. C., Yeast regulatory gene *PPR1*. I. Nucleotide sequence, restriction map and codon usage, 239
— (see also Liljelund, P.), 251
Keller, W. (see also Sergeant, A.), 577
Khandekar, P., Saidapet, C., Krauskopf, M., Zarraga, A.-M., Lin, W.-L., Mendola, C. & Siddiqui, M. A. Q., Co-ordinate control of gene expression, Muscle-specific 7 S RNA contains sequences homologous to 3′-untranslated regions of myosin genes and repetitive DNA, 417
Khalili, K. & Weinmann, R., Actin mRNAs in HeLa cells. Stabilization after adenovirus infection, 1007
King, J. (see also Bryant, J. L. Jr), 837
Kirov, N., Djondjurov, L. & Tsanev, R., Nuclear matrix and transcriptional activity of the mouse α-globin gene, 601
Knight, P. (see also Trinick, J.), 331
Kole, A. (see also Malan, T. P.), 881
Koop, B. F. (see also Goodman, M.), 803
Krause, M. (see also Landel, C. P.), 497
Krauskopf, M. (see also Khandekar, P.), 417
Krieg, P., Strachan, R , Wallis, E., Tabe, L. & Colman, A., Efficient expression of cloned complementary DNAs for secretory proteins after injection into *Xemopus* oocytes, 615
— (see also Tabe, L.), 645

L1 RNA maturation is interfered with by a specific splicing block; expression of two *Xenopus laevis* ribosomal protein genes in injected frog ooxytes, 987
La Mar, G. N. (see also Lecomte, J.), 357
lac operon transcription initiation activation of the P1 promoter, mechanism of CRP-cAMP activation, 881
— repressor, *Escherichia coli*, amino acid replacements in the DNA binding domain, 205
Lacroute, F. (see also Liljelund, P.), 251
λ (see also Bacteriophage), 41, 283, 865, 1053
Landel, C. P., Krause, M., Waterston, R. H. & Hirsh, D., DNA rearrangements of the actin gene cluster in *Caenorhabditis elegans* accompany reversion of three muscle mutants, 497
Lane, D., Hill, D., Caughey, P. & Gunn, P., The mini-F primary origin. Sequence analysis and multiple activities, 267
— T. R. (see also Waterston, R. H.), 473
Lazdunski, C. (see also Varenne, S.), 549
Lebioda, L. & Brewer, J. M., Crystallization and preliminary crystallographic data for a tetragonal form of yeast enolase, 213
Leclerc, J. E., Istock, N. L., Saran, B. R. & Allen, R. Jr, Sequence analysis of ultraviolet-induced mutations in M13*lacZ* hybrid phage DNA, 217
Lecomte, J., La Mar, G. N., Winterhalter, K. H. & Smit, J. D. G., Proton nuclear magnetic resonance investigation of the heme cavity structure of liver fluke (*Dicrocoelium dendriticum*) methemoglobin, 357
Levis, R. (see also O'Hare, K.), 437
Liljelund, P., Losson, R., Kammerer, B. & Lacroute, F., Yeast regulatory gene *PPR1*. II. Chromosomal localization, meiotic map, suppressibility, dominance/recessivity and dosage effect, 251
Lilley, D. M. J. & Hallam, L. R., Thermodynamics of the ColE1 cruciform. Comparison between probing and topological experiments using single topoisomers, 179
Lin, W.-L. & Pederson, T., Ribonucleoprotein organization of eukaryotic RNA. XXXI. Structure of the U1 small nuclear ribonucleoprotein, 947
— (see also Khandekar, P.), 417
Linker histone composition variations control RNA polymerase binding to chromatin, 131
Liver fluke methemoglobin, proton n.m.r. investigation of the heme cavity structure, 357
Lloubes, R. (see also Varenne, S.), 549
Losson, R. (see also Liljelund, P.), 251
Loubresse, N. G. (see also Adoutte, A.), 1065
Luo, M., Arnold, E., Erickson, J. W., Rossmann, M. G., Boege, U. & Scraba, D. G., Picornaviruses of two different genera have similar structures, 703
Lustig, A. J. & Peters, T. D., Long poly(A) tracts in the human genome are associated with the *Alu* family of repeated elements, 753

Mcclure, W. R. (see also Malan, T. P.), 881
McDonald, M. J. (see also Moo-Penn, W. F.), 1119
Mahoney, M. E. (see also Place, N.), 865
Malan, T. P., Kole, A., Buc, H. & McClure, W. R., Mechanism of CRP-cAMP activation of *lac* operon transcription initiation activation of the P1 promoter, 881
Mammals, the long evolutionary history of the η-globin gene in the β-globin gene family, 803
Matrix, nuclear, and transcriptional activity of the mouse α-globin gene, 601
Matsuoka, A. (see also Takagi, T.), 1179
Meiotic map, yeast regulatory gene *PPR1*, 251
Membrane protein complex, X-ray structure analysis. Electron density map at 3 Å resolution and a model of the chromophores of the photosynthetic reaction centre from *Rhodopseudomonas viridis*, 385
Mendola, C. (see also Khandekar, P.), 417
Messenger RNAs, actin, in HeLa cells. Stabilization after adenovirus infection, 1007
Methemoglobin, liver fluke, a proton n.m.r. investigation of the heme cavity structure, 357
Michel, H. (see also Deisenhofer, J.), 385
Miki, K. (see also Deisenhofer, J.), 385
Miller, B. L. (see also Gwynne, B. L.), 91
—, J. H., Genetic studies of the *lac* repressor. XII. Amino acid replacements in the DNA binding domain of the *Escherichia coli lac* repressor, 205
—, K. Y. (see also Gwynne, D. I.), 91

Mini-F primary origin. Sequence analysis and multiple activities, 267
M13*lacZ* hybrid phage DNA, sequence analysis of ultraviolet-induced mutations, 217
Monaco, H. L. (see also Zanette, D.), 1185
Monoclonal antibody Fabs, crystalline, complexed to hen egg white lysozyme, 761
Moo-Penn, W. F., Jue, D. L., Johnson, M. H., McDonald, M. J., Turci, S. M., Shih, T.-B., Jones, R. T., Therrell, B. L. Jr & Arnone, A., Structural and functional studies of hemoglobin Wayne: an elongated α-chain variant, 1119
Motile domains in the subfragment-2 region of myosin, an enzyme-probe study, 667
Mouse α-globin gene, nuclear matrix and transcriptional activity, 601
Murphy, C. (see also O'Hare, K.), 437
Muscle mutants, three, reversion is accompanied by DNA rearrangements of the actin gene cluster in *Caenorhabditis elegans*, 497
—, skeletal, vertebrate, the thin filament co-operatively activates as a unit, 379
Muscle structure dominant mutations in *Caenorhabditis elegans* that map near the actin gene cluster, 473
—-specific 7 S RNA contains sequences homologous to 3′-untranslated regions of myosin genes and repetitive DNA, co-ordinate control of gene expression, 417
Mutant ovalbumins and ovalbumin–globin fusion proteins, segregation in *Xenopus* oocytes, 645
Mutants, muscle, three, reversion is accompanied by DNA rearrangements of the actin gene cluster in *Caenorhabditis elegans*, 497
Mutations, dominant, affecting muscle structure in *Caenorhabditis elegans* that map near the actin gene cluster, 473
— that alter the DNA binding site for the bacteriophage lambda *c*II protein and affect the translation efficiency of the *c*II gene, 865
Mutations, ultraviolet-induced, in M13*lacZ* hybrid phage DNA, sequence analysis, 217
Myoglobins, *Aplysia*, with an unusual amino acid sequence, 1179
Myosin, an enzyme-probe study of motile domains in the subfragment-2 region, 667
— genes and repetitive DNA, muscle-specific 7 S RNA contains sequences homologous to 3′-untranslated regions; co-ordinate control of gene expression, 417
Myslinski, E., Branlant, C., Wieben, E. D. & Pederson, T., The small nuclear RNAs of *Drosophila*, 927

N antitermination function of phage lambda, and the sequence or position of the *nusA* recognition site, 1053
Neutron and X-ray refinement of crystal form II structure of bovine pancreatic trypsin inhibitor, 301
Noller, H. F. (see also Van Stolk, B. J.), 151
Nuclear magnetic resonance assignments and structural comparison of wild-type and mutant λ OR3 operator DNA; sequence-specific recognition of DNA, 41
— — — determination of polypeptide secondary structure by observation of short proton–proton distances, 715
— — — investigation of the heme cavity structure of liver fluke methemoglobin, 357
— matrix and transcriptional activity of the mouse α-globin gene, 601
Nucleotide sequence analysis of the β-type globin pseudogene from humans, gorilla and chimpanzee, 767
— — of bacteriophage T4 gene *23* and the amino acid sequence of its product, 399
— and transcription of the phenylalanine and tyrosine operons of *Escherichia coli* K12, 1023
— — of yeast regulatory gene *PPR1*, 239
nusA recognition site: alteration in its sequence or position relative to upstream translation interferes with the action of the N antitermination function of phage lambda, 1053

O'Hare, K., Murphy, C., Levis, R. & Rubin, G. M., DNA sequence of the *white* locus of *Drosophila melanogaster*, 437
Olson, E. R., Tomich, C.-S. C. & Friedman, D. I., The *nusA* recognition site. Alteration in its sequence or position relative to upstream translation interferes with the action of the N antitermination function of phage lambda, 1053
Oocytes, frog, injected, and the expression of two *Xenopus laevis* ribosomal protein genes; a specific block interferes with the L1 RNA maturation, 987
— *Xenopus*, efficient expression of cloned complementary DNAs for secretory proteins after injection, 615
— —, segregation of mutant ovalbumins and ovalbumin–globin fusion proteins, 645

Ovalbumins, mutant, and ovalbumin–globin fusion proteins, segregation in *Xenopus* oocytes, 645
Oxygen affinity of bovine haemoglobin; relevance of electrostatic and hydrophobic interactions, 1157

P_C and P_{BAD} *Escherichia coli ara* promoters, deletion analysis, 201
P1 promoter, mechanism of CRP-cAMP activation of *lac* operon transcription initiation activation, 881
P22 (see also Bacteriophage), 837
Padlan, E. A. (see also Silverton, E. W.), 761
Palfreyman, J. W. (see also Campbell, M. E. M.), 1
Palindromic and non-palindromic sequences, role in arresting DNA synthesis *in vitro* and *in vivo*, 961
Paramecium, proteolytic cleavage and maturation of the crystalline secretion products, 1065
Pardi, A., Billeter, M. & Wüthrich, K., Calibration of the angular dependence of the amide proton–C^α proton coupling constants, $^3J_{HN\alpha}$, in a globular protein. Use of $^3J_{HN\alpha}$ for identification of helical secondary structure, 741
Parge, H. E., Schneider, M., Hahn, U., Saenger, W., Altschmied, L. & Hillen, W., Crystallization of and preliminary X-ray diffraction data for TET-repressor and the TET-repressor–tetracycline complex, 1189
Parker, M. L., Christensen, A. C., Boosman, A., Stockard, J., Young, E. T. & Doermann, A. H., Nucleotide sequence of bacteriophage T4 gene *23* and the amino acid sequence of its product, 399
Pederson, T. (see also Lin, W.-L.), 947
— (see also Myslinski, E.), 927
— (see also Sass, H.), 911
Pero, J. (see also Costanzo, M.), 533
Petes, T. D. (see also Lustig, A. J.), 753
Phenylalanine and tyrosine operons of *Escherichia coli* K12, nucleotide sequence and transcription, 1023
Photoinactivation, acridine-sensitized, of bacteriophage P22; DNA injection proteins are targets, 837
Picornaviruses of two different genera have similar structures, 703
Pierandrei-Amaldi, P. (see also Bozzoni, I.), 987, 1173
Place, N., Fien, K., Mahoney, M. E., Wulff, D. L., Ho, Y-S., Debouck, C., Rosenberg, M., Shih, M.-C. & Gussin, G. N., Mutations that alter the DNA binding site for the bacteriophage lambda *c*II protein and affect the translation efficiency of the *c*II gene, 865
Poly(A) long tracts in the human genome are associated with the *Alu* family of repeated elements, 753
Polypeptide chains, nascent, effect of tRNA availability on the rate of elongation; translation is a non-uniform process, 549
Polypeptide secondary structure determination by n.m.r. observation of short proton–proton distances, 715
Polytene chromosomes, transcription-dependent localization of U1 and U2 small nuclear ribonucleoproteins at major sites of gene activity, 911
Potter, M. (see also Silverton, E. W.), 761
PPR1 yeast regulatory gene; chromosomal localization, meiotic map, suppressibility, dominance/recessivity and dosage effect, 251
PPR1 yeast regulatory gene; nucleotide sequence, restriction map and codon usage, 239
Preston, C. M. (see also Campbell, M. E. M.), 1
Primate $\psi\beta 1$ gene. An ancient β-globin pseudogene, 785
Probing and topological experiments using single topoisomers, a comparison; thermodynamics of the ColE1 cruciform, 179
Prohead binding, a functional domain of bacteriophage λ terminase, 283
Proline peptide bonds, and short proton–proton distances in polypeptide chains, 734
Proline/alanine alternating sequence of βB1 subunit originates from a repetitive DNA sequence. Bovine β-crystalline complementary DNA clones, 457
Protein, riboflavin-binding, hen eggwhite; crystallization, 1185
Proteolytic cleavage and maturation of the crystalline secretion products of *Paramecium*, 1065
Proton exchange, amide, used to monitor the formation of a stable α-helix by residues 3 to 13 during folding of ribonuclease S, 1141
— n.m.r. investigation of the heme cavity structure of liver fluke (*Dicrocoelium dendriticum*) methemoglobin, 357
Proton–C^α proton coupling constants, amide, $^3J_{HN\alpha}$, in a globular protein, calibration of the angular dependence; use for identification of helical secondary structure, 741
Proton–proton distances, short, in polypeptide chains containing glycine, proline or *cis* peptide bonds, 734

———, —, and polypeptide secondary structure determination by n.m.r. observation, 715
Pseudogene, β-globin, ancient; the primate ψβ1 gene, 785
—, β-type globin, from human, gorilla and chimpanzee, isolation and nucleotide sequence analysis, 767
ψβ1 gene, primate; an ancient β-globin pseudogene, 785

Quax, W. (see also Quax-Jeuken, Y.), 457
Quax-Jeuken, Y., Janssen, C., Quax, W., Van Den Heuvel, R. & Bloemendal, H., Bovine β-crystallin complementary DNA clones. Alternating proline/alanine sequence of βB1 subunit originates from a repetitive DNA sequence, 457

Recombinational hotspot activity of Chi-like sequences, 371
Reid, B. R. (see also Wemmer, D. E.), 41
Regulatory subunits of *Bacillus subtilis* RNA polymerase, location and primary structure of encoding genes; bacteriophage SPO1 genes *33* and *34*, 533
Repeated elements, *Alu* family, are associated with long poly(A) tracts in the human genome, 753
Restriction map, yeast regulatory gene *PPR1*, 239
Rhodopseudomonas viridis, electron density map at 3 Å resolution and a model of the chromophores of the photosynthetic reaction centre, 385
Riboflavin-binding protein, hen eggwhite; crystallization, 1185
Ribonuclease S folding, and amide proton exchange used to monitor the formation of a stable α-helix by residues 3 to 13, 1141
RNA, eukaryotic; structure of the U1 small nuclear ribonucleoprotein, 947
— polymerase, *Bacillus subtilis*, location and primary structure of genes encoding regulatory subunits; bacteriophage SPO1 genes *33* and *34*, 533
— — binding to chromatin, control by variations in linker histone composition, 131
—, 7 S, muscle-specific, contains sequences homologous to 3′-untranslated regions of myosin genes and repetitive DNA; co-ordinate control of gene expression, 417
RNAs, ribosomal protein, *Xenopus laevis*, splicing is inhibited *in vivo* by antisera to ribonucleoproteins containing U1 small nuclear RNA, 1173
RNA, ribosomal, 16 S, chemical probing of conformation using diethylpyrocarbonate, 151
RNAs, small nuclear, of *Drosophila*, 927
Ribonucleoprotein organization of eukaryotic RNA. Structure of the U1 small nuclear ribonucleoprotein, 947
Ribonucleoproteins containing U1 small nuclear RNA, antisera inhibit *in vivo* the splicing of *Xenopus laevis* ribosomal protein RNAs, 1173
Ribosomal DNA in *Dictyostelium discoideum*, site-specific phasing in the chromatin, 73
— RNA, 16 S, chemical probing of conformation using diethylpyrocarbonate, 151
— protein genes, two, *Xenopus laevis*, expression in injected frog oocytes; a specific block interferes with the L1 RNA maturation, 987
— — RNAs, *Xenopus laevis*, splicing is inhibited *in vivo* by antisera to ribonucleoproteins containing U1 small nuclear RNA, 1173
Rosenberg, M. (see also Place, N.), 865
Rossmann, M. G. (see also Luo, M.), 703
Rubin, G. M. (see also O'Hare, K.), 437

Saenger, W. (see also Parge, H. E.), 1189
Saidapet, C. (see also Khandekar, P.), 417
Saran, B. R. (see also Leclerc, J. E.), 217
Sass, H. & Pederson, T., Transcription-dependent localization of U1 and U2 small nuclear ribonucleoproteins at major sites of gene activity in polytene chromosomes, 911
Schachat, F. H. (see also Brandt, P. W.), 379
Schleif, R. (see also Dunn, T. M.), 201
— (see also Hahn, S.), 61
Schmidt, T. (see also Alonso, A.), 825
Schneider, M. (see also Parge, H. E.), 1189
Scraba, D. G. (see also Luo, M.), 703
Second large SV40 T-antigen exon contains the information for maximal cell transformation, 111
Secretion products, crystalline, of *Paramecium*; proteolytic cleavage and maturation, 1065
Segregation of mutant ovalbumins and ovalbumin–globin fusion proteins in *Xenopus* oocytes, 645

Sequence, alternating proline/alanine, of βB1 subunit originates from a repetitive DNA sequence. Bovine β-crystallin complementary DNA clones, 457
—, amino acid, unusual, in *Aplysia* myoglobins, 1179
— analysis and multiple activities; mini-F primary origin, 267
— — of ultraviolet-induced mutations in M13*lacZ* hybrid phage DNA, 217
—, DNA, of the *white* locus of *Drosophila melanogaster*, 437
—, nucleotide, analysis of the β-type globin pseudogene from human, gorilla and chimpanzee, 767
—, —, of bacteriophage T4 gene *23* and the amino acid sequence of its product, 399
—, , and transcription of the phenylalanine and tyrosine operons of *Escherichia coli* K12, 1023
—, —, of yeast regulatory gene *PPR1*, 239
Sequences, Chi-like, recombinational hotspot activity, 371
, DNA, herpes simplex virus, which encode a *trans*-acting polypeptide responsible for stimulation of immediate early transcription, 1
— homologous to 3′-untranslated regions of myosin genes and repetitive DNA are contained in muscle-specific 7 S RNA; co-ordinate control of gene expression, 417
, palindromic and non-palindromic, role in arresting DNA synthesis *in vitro* and *in vivo*, 961
Sequence-specific recognition of DNA. Nuclear magnetic resonance assignments and structural comparison of wild-type and mutant λ OR3 operator DNA, 41
Sergeant, A., Bohmann, D., Zentgraf, H., Weiher, H. & Keller, W., A transcription enhancer acts *in vitro* over distances of hundreds of base-pairs on both circular and linear templates but not on chromatin-reconstituted DNA, 577
Shepherd, B. M. & Finnegan, D. J., Structure of circular copies of the *412* transposable element present in *Drosophila melanogaster* tissue culture cells, and isolation of a free *412* long terminal repeat, 21
Shikama, K. (see also Takagi, T.), 1179
Shih, M.-C. (see also Place, N.), 865
—, T.-B. (see also Moo-Penn, W. F.), 1119
Siddiqui, M. A. Q. (see also Khandekar, P.), 417
Siegele, D. A. (see also Frackman, S.), 283
Silverton, E. W., Padlan, E. A., Davies, D. R., Smith-Gill, S. & Potter, M., Crystalline monoclonal antibody Fabs complexed to hen egg white lysozyme, 761
Simian virus 40 second large T-antigen exon contains the information for maximal cell transformation, 111
Site-specific phasing in the chromatin of the rDNA in *Dictyostelium discoideum*, 73
Sjölin, L. (see also Wlodawer, A.), 301
Slightom, J. L. (see also Chang, L.-Y. E.), 767
— (see also Goodman, M.), 803
Small nuclear ribonucleoprotein, U1, structure, 947
— — ribonucleoproteins, U1 and U2, transcription-dependent localization at major sites of gene activity in polytene chromosomes, 911
— — RNA genes, U1, *Drosophila melanogaster*, 825
— -, U1, contained in ribonucleoproteins, antisera inhibit *in vivo* the splicing of *Xenopus laevis* ribosomal protein RNAs, 1173
Small nuclear RNAs of *Drosophila*, 927
Smit, J. D. G. (see also Lecomte, J.), 357
Smith, G. R. (see also Cheng, K. C.), 371
Smith-Gill, S. (see also Silverton, E. W.), 761
Spadon, P. (see also Zanette, D.), 1185
Splicing of *Xenopus laevis* ribosomal protein RNAs is inhibited *in vivo* by antisera to ribonucleoproteins containing U1 small nuclear RNA, 1173
SPO1 (see also Bacteriophage), 533
SpoC1 gene cluster from *Aspergillus nidulans*, structure and regulated expression, 91
Stockard, J. (see also Parker, M. L.), 399
Strachan, R. see also Krieg, P.), 165
— (see also Tabe, L.), 645
Structural comparison of purified dynein proteins with *in situ* dynein arms, 1083
— and functional studies of hemoglobin Wayne: an elongated α-chain variant, 1119
— of bovine pancreatic trypsin inhibitor. Results of joint neutron and X-ray refinement of crystal form II, 301
— — circular copies of the *412* transposable element present in *Drosophila melanogaster* tissue culture cells, and isolation of a free *412* long terminal repeat, 21

, helical, secondary, identified with amide proton $-C^{\alpha}$ proton coupling constants, $^{3}J_{HN\alpha}$, 741
—, heme cavity, of liver fluke methemoglobin, a proton n.m.r. investigation, 357
, muscle, and dominant mutations in *Caenorhabditis elegans* that map near the actin gene cluster, 473
—, polypeptide, secondary, determined by n.m.r. observation of short proton–proton distances, 715
, primary, and location of genes encoding regulatory subunits of *Bacillus subtilis* RNA polymerase; bacteriophage SPO1 genes *33* and *34*, 533
— and regulated expression of the SpoC1 gene cluster from *Aspergillus nidulans*, 91
— of the U1 small nuclear ribonucleoprotein, 947
-, X-ray, analysis of a membrane protein complex. Electron density map at 3 Å resolution and a model of the chromophores of the photosynthetic reaction centre from *Rhodopseudomonas viridis*, 385
Structures, similar, for picornaviruses of two different genera, 703
Subfragment-2 region of myosin, an enzyme-probe study of motile domains, 667
Suppressibility, yeast regulatory gene *PPR1*, 251

T4 (see also Bacteriophage), 399
Tabe, L., Krieg, P., Strachan, R., Jackson, D., Wallis, E. & Colman, A., Segregation of mutant ovalbumins and ovalbumin-globin fusion proteins in *Xenopus* oocytes, 645
— (see also Krieg, P.), 615
Takagi, T., Iida, S., Matsuoka, A. & Shikama, K., *Aplysia* myoglobins with an unusual amino acid sequence, 1179
T-antigen exon, second large SV40, contains the information for maximal cell transformation, 111
Terminase, bacteriophage λ, a functional domain for prohead binding, 283
Tetracycline complex of TET-repressor, crystallization and preliminary X-ray diffraction data, 1189
TET-repressor and the TET-repressor–tetracycline complex, crystallization and preliminary X-ray diffraction data, 1189
Thermodynamics of the ColE1 cruciform. Comparisons between probing and topological experiments using single topoisomers, 179
Therrell, B. L. Jr (see also Moo-Penn, W. F.), 1119
Timberlake, W. E. (see also Gwynne, D. I.), 91
Titin, native, purification and properties, 331
Tomich, C.-S. C. (see also Olson, E. R.), 1053
Topoisomerase, eukaryotic type 1, breakage of single-stranded DNA occurs only at regions with the potential for base-pairing, 515
Topological and probing experiments using single topoisomers, a comparison; thermodynamics of the ColE1 cruciform, 179
Trans-acting polypeptide responsible for stimulation of immediate early transcription, identification of encoding herpes simplex virus DNA sequences, 1
Transcription antitermination function of phage lambda, alteration in the sequence or position of the *nusA* recognition site interferes with the action, 1053
enhancer acts *in vitro* over distances of hundreds of base-pairs on both circular and linear templates but not on chromatin-reconstituted DNA, 577
—, immediate early; identification of herpes simplex virus DNA sequences which encode a *trans*-acting polypeptide responsible for stimulation, 1
initiation activation, *lac* operon, of the P1 promoter, mechanism of CRP-cAMP activation, 881
— of the phenylalanine and tyrosine operons of *Escherichia coli* K12, 1023
Transcription-dependent localization of U1 and U2 small nuclear ribonucleoproteins at major sites of gene activity in polytene chromosomes, 911
Transcriptional activity and nuclear matrix of the mouse α-globin gene, 601
Transfer RNA availability, effect on the rate of elongation of nascent polypeptide chains. Translation is a non-uniform process, 549
Translation efficiency of the *c*II gene is affected by mutations that alter the DNA binding site for the bacteriophage lambda *c*II protein, 865
— is a non-uniform process. Effect of tRNA availability on the rate of elongation of nascent polypeptide chains, 549
Trinick, J., Knight, P. & Whiting, A., Purification and properties of native titin, 331
Trypsin inhibitor, pancreatic, bovine; structure; joint neutron and X-ray refinement of crystal form II, 301
Tsanev, R. (see also Kirov, N.), 601

Turci, S. M. (see also Moo-Penn, W. F.), 1119
Tyrosine and phenylalanine operons of *Escherichia coli* K12, nucleotide sequence and transcription, 1023

U1 small nuclear ribonucleoprotein structure, 947
— — — RNA-containing ribonucleoproteins, antisera inhibit *in vivo* the splicing of *Xenopus laevis* ribosomal protein RNAs, 1173
— snRNA genes of *Drosophila melanogaster*, 825
— and U2 small nuclear ribonucleoproteins, transcription-dependent localization at major sites of gene activity in polytene chromosome, 911
Ueno, H. & Harrington, W. F., An enzyme-probe study of motile domains in the subfragment-2 region of myosin, 667
Ultraviolet-induced mutations in M13*lacZ* hybrid phage DNA, sequence analysis, 217
Upstream repression and CRP stimulation of the *Escherichia coli* L-arabinose operon, 61

Van Den Heuvel, R. (see also Quax-Jeuken, Y.), 457
Van Stolk, B. J. & Noller, H. F., Chemical probing of conformation in large RNA molecules. Analysis of 16 S ribosomal RNA using diethylpyrocarbonate, 151
Varenne, S., Buc, J., Lloubes, R. & Lazdunski, C., Translation is a non-uniform process. Effect of tRNA availability on the rate of elongation of nascent polypeptide chains, 549
Vertebrate skeletal muscle, the thin filament co-operatively activates as a unit, 379
Vitrano, E., Cupane, A. & Cordone, L., Oxygen affinity of bovine haemoglobin, Relevance of electrostatic and hydrophobic interactions, 1157

Wallis, E. (see also Krieg, P.), 615
— (see also Tabe, L.), 645
Walter, J. (see also Wlodawer, A.), 301
Waterston, R. H., Hirsh, D. & Lane, T. R., Dominant mutations affecting muscle structure in *Caenorhabditis elegans* that map near the actin gene cluster, 473
Waterston, R. H. (see also Landel, C. P.), 497
Wayne, hemoglobin, and elongated α-chain variant; structural and functional studies, 1119
Weaver, D. T. & DePamphilis, M. L., The role of palindromic and non-palindromic sequences in arresting DNA synthesis *in vitro* and *in vivo*, 961
Weiher, H. (see also Sergeant, A.), 577
Weinmann, R. (see also Khalili, K.), 1007
Weiss, M. L. (see also Goodman, M.), 803
Weiss, M. L. (see also Harris, S.), 785
Wemmer, D. E., Chou, S.-H. & Reid, B. R., Sequence-specific recognition of DNA. Nuclear magnetic resonance assignments and structural comparison of wild-type and mutant λ OR3 operator DNA, 41
white locus DNA sequence, *Drosophila melanogaster*, 437
Whiting, A. (see also Trinick, J.), 331
Wieben, E. D. (see also Myslinski, E.), 927
Winterhalter, K. H. (see also Lecomte, J.), 357
Wlodawer, A., Walter, J., Huber, R. & Sjölin, L., Structure of bovine pancreatic trypsin inhibitor. Results of joint neutron and X-ray refinement of crystal form II, 301
Wulff, D. L. (see also Place, N.), 865
Wüthrich, K., Billeter, M. & Braun, W., Polypeptide secondary structure determination by nuclear magnetic resonance observation of short proton–proton distances, 715
— (see also Pardi, A.), 741

Xenopus laevis, expression of two ribosomal protein genes in injected frog oocytes. A specific splicing block interferes with the L1 RNA maturation, 987
— — ribosomal protein RNA splicing is inhibited *in vivo* by antisera to ribonucleoproteins containing U1 small nuclear RNA, 1173
— oocytes, efficient expression of cloned complementary DNAs for secretory proteins after injection, 615
— —, segregation of mutant ovalbumins and ovalbumin–globin fusion proteins, 645

X-ray and neutron refinement of crystal form II structure of bovine pancreatic trypsin inhibitor, 301
— preliminary diffraction data for TET-repressor and the TET-repressor–tetracycline complex, 1189
— structure analysis of a membrane protein complex. Electron density map at 3 Å resolution and a model of the chromophores of the photosynthetic reaction centre from *Rhodopseudomonas viridis*, 385

Yeast enolase, crystallization and preliminary crystallographic data for a tetragonal form, 213
— regulatory gene *PPR1*. Chromosomal localization, meiotic map, suppressibility, dominance/recessivity and dosage effect, 251
— — — —. Nucleotide sequence, restriction map and codon usage, 239
Young, E. T. (see also Parker, M. L.), 399

Zanette, D., Monaco, H. L., Zanotti, G. & Spadon, P., Crystallization of hen eggwhite riboflavin-binding protein, 1185
Zanotti, G. (see also Zanette, D.), 1185
Zarraga, A.-M. (see also Khandekar, P.), 417
Zentgraf, H. (see also Sergeant, A.), 577